

| | |
|-------------|--|
| Seat No. | |
|-------------|--|

[5057]-251**S.E. (Comp/IT) (Second Semester) EXAMINATION, 2016****ENGINEERING MATHEMATICS-III****(2012 PATTERN)****Time : Two Hours****Maximum Marks : 50**

N.B. :- (i) Answer Q. No. 1 or Q. No. 2, Q. No. 3 or Q. No. 4,
Q. No. 5 or Q. No. 6, Q. No. 7 or Q. No. 8.

(ii) Neat diagrams must be drawn wherever necessary.

(iii) Figures to the right indicate full marks.

(iv) Use of logarithmic tables, slide rule, Mollier charts, electronic pocket calculator and steam tables is allowed.

(v) Assume suitable data, if necessary.

1. (a) Solve any two : [8]

(i) $(D^2 - 1)y = \cos x \cosh x + 3^x$

(ii) $(D^2 + 3D + 2)y = e^{e^x}$

(iii) $(2x+3)^2 \frac{d^2y}{dx^2} + (2x+3) \frac{dy}{dx} - 2y = 24x^2$.

(b) Find the Fourier transform of [4]

$$f(x) = \begin{cases} 1-x^2, & |x| \leq 1 \\ 0, & |x| > 1 \end{cases}$$

Hence evaluate $\int_0^{\infty} \left(\frac{x \cos x - \sin x}{x^3} \right) \cos \frac{x}{2} dx$.

Or

2. (a) An e.m.f. $E \sin pt$ is applied at $t = 0$ to a circuit containing a condenser C and inductance L in series, the current I satisfies the equation $L \frac{di}{dt} + \frac{1}{C} \int i dt = E \sin pt$, where

$$i = -\frac{dq}{dt}, \text{ if } p^2 = \frac{1}{LC} \text{ and}$$

initially the current and the charge are zero, find current at any time t . [4]

- (b) Find the inverse z -transform (any one) : [4]

(i) $F(z) = \frac{1}{(z-3)(z-4)}, |z| < 3$

- (ii) Find inverse z -transform of $F(z) = \frac{z^2}{z^2+1}$ using inversion integral method.

- (c) Solve the following difference equation to find $f(k)$. [4]

$$12f(k+2) - 7f(k+1) + f(k) = 0$$

$$k \geq 0, f(0) = 0, f(1) = 3.$$

3. (a) The first four moments of a distribution about 30.2 are 0.255, 6.222, 30.211 and 400.25. Calculate the first four moments about the mean. Also calculate coefficient of skewness. [4]

- (b) Suppose heights of students follows normal distribution with mean 190 cm and variance 80 cm^2 . In a school of 1000 students, how many would you expect to be above 200 cm tall ?

(Given that : $A_1(z > 1.1180) = 0.13136$). [4]

- (c) Find the directional derivative of $\phi = e^{2x} \cos(yz)$ at $(0, 0, 0)$ in the direction of tangent to the curve $x = a \sin t, y = a \cos t, z = at$, at $t = \frac{\pi}{4}$. [4]

Or

4. (a) Prove the following (any one) : [4]

(i) $\nabla \cdot \left(\frac{\vec{a} \times \vec{r}}{r} \right) = 0$

(ii) $\nabla^4 (r^2 \log r) = \frac{6}{r^2}$.

- (b) Show that vector field given by $\vec{F} = (y^2 \cos x + z^2)\vec{i} + (2y \sin x)\vec{j} + (2xz)\vec{k}$ is conservative and find scalar field ϕ such that $\vec{F} = \nabla\phi$. [4]

- (c) If $\sum x_i = 30, \sum y_i = 40, \sum x_i^2 = 220, n = 5, \sum y_i^2 = 340$ and $\sum x_i y_i = 214$, then obtain the regression lines for this data. [4]

5. (a) Evaluate the integral $\int \vec{F} \cdot d\vec{r}$, where

$$\vec{F} = (y \sin z - \sin x)\vec{i} + (x \sin z + 2yz)\vec{j} + (x \cos z + y^2)\vec{k}$$

from the point $(0, 0, 0)$ to $\left(\frac{\pi}{2}, 1, \frac{\pi}{2}\right)$. Is \vec{F} conservative ?

[5]

- (b) Using divergence theorem, evaluate $\iint_S \vec{F} \cdot \hat{n} \, ds$, where $\vec{F} = x\hat{i} - y\hat{j} + (z^2 - 1)\hat{k}$ and S is the total surfaces of the cylinder bounded by $z = 0$, $z = 1$ and $x^2 + y^2 = 4$. [4]
- (c) Use Stokes' theorem to evaluate $\iint_S (\nabla \times \vec{F}) \cdot \hat{n} \, ds$, where $\vec{F} = yi + (x - 2xz)j - xyk$ and S is the surface of the sphere $x^2 + y^2 + z^2 = a^2$, above the xy plane. [4]

Or

6. (a) Evaluate the integral $\int_C \vec{F} \cdot d\vec{r}$, where $\vec{F} = [e^x y + \sin y]\hat{i} + [e^x + x(1 + \cos y)]\hat{j}$ where C is the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, z = 0$. [4]
- (b) Evaluate $\iint_S (yz\hat{i} + zx\hat{j} + xy\hat{k}) \cdot \hat{n} \, dS$, where S is the curved surface of the cone $x^2 + y^2 = z^2, z = 4$. [5]
- (c) If $\vec{E} = \nabla\phi$ and $\nabla^2\phi = -4\pi\rho$, prove that :

$$\iint_S \vec{E} \cdot d\vec{s} = -4\pi \iiint_V \rho \, dV. \quad [4]$$

7. (a) Find the harmonic conjugate of $v = e^x \sin y$ such that $f(z) = u + iv$ is analytic. Find $f(z)$ in terms of z . [4]
- (b) Using Cauchy's Integral formula evaluate $\oint_C \frac{3z^3 + 5z + 2}{(z-2)^2} dz$ where C is $\frac{x^2}{9} + \frac{y^2}{25} = 1$. [5]

- (c) Find the map of the strip $x > 0$, $0 < y < 4$ under the transformation $w = iz + 2$. [4]

Or

8. (a) Show that analytic function with constant amplitude is constant. [4]

(b) Evaluate $\oint_C \frac{z-3}{z^2+2z+5} dz$, where 'C' is $|z| = 1$. [5]

- (c) Find the bilinear transformation which maps the points $z = 0, 1, 2$ onto the points $w = 1, \frac{1}{2}, \frac{1}{3}$ respectively. [4]

stupidstupid.com

