

Total No of Questions: [8]

SEAT NO.

[Total No. of Pages : 3]

S.E. 2012 (Electronics / E &TC)

Engineering Mathematics – III

(Semester - I)

Time: 2 Hours

Max. Marks : 50

Instructions to the candidates:

- 1) Answers Q1 or Q2, Q3 or Q4, Q5 or Q6, Q7 or Q8.
- 2) Neat diagrams must be drawn wherever necessary.
- 3) Figures to the right side indicate full marks.
- 4) Use of Calculator is allowed.
- 5) Assume Suitable data if necessary

- Q1) a) Solve (any two) [8]
- i) $(D^2-1)y = x \sin x + (1+x^2)e^x$
 - ii) $d^2y/dx^2 + y = \operatorname{cosec} x$ (by variation of parameters)
 - iii) $x^2 d^2y/dx^2 - 4x dy/dx + 6y = x^5$
- b) Find Fourier cosine transform of the function [4]
- $$f(x) = \begin{cases} \cos x & 0 < x < a \\ 0 & x > a \end{cases}$$

OR

- Q2) a) A resistance of 50 ohms, an inductor of 2 henries and farad capacitor [4]
are all in series with an e.m.f. of 40 volts . Find the instantaneous
change and current after the switch is closed at $t=0$, assuming that at
that time the charge on the capacitor is 4 coulomb.
- b) Solve (any one) [4]
- i) Find z transform for $f(k) = (1/3)^{|k|}$
 - ii) Find inverse z transform of $[Z^2 / (Z-1/4) (Z-1/5)]$ for $|Z| < 1/5$
 - c) Solve $f(k+2) + 3 f(k+1) + 2 f(k) = 0$ [4]
Given $f(0) = 0$, $f(1)=1$

- Q3) a) Solve the following differential equation to get $y(0.1)$ [4]
 $dy/dx = x-y^2$, $y(0) = 1$
by using Runge- Kutta fourth order method. ($h=0.1$)
- b) Find Lagrange's long interpolating polynomial passing through set of [4]
points

x	0	1	2
y	2	3	6

Use it to find y at x = 1.5 and find $\int_0^2 y dx$.

- c) Find the directional derivative of $\phi = 3 \log (x+y+z)$ at (1,1,1) in the direction of tangent to the curve $x = b \sin t, y = b \cos t, z = bt$ at $t=0$ [4]

OR

- Q4) a) Show that (any one) [4]

i) $\nabla^2[\nabla \cdot (\vec{r}/r^2)] = 2/r^4$

ii) $\nabla(a^{-1} \cdot \vec{r}^{-1} / r^3) = a^{-1} / r^3 - 3(a^{-1} \cdot \vec{r}^{-1}) / r^5$

- b) If ϕ, ψ satisfy Laplace equation then prove that the vector $(\phi \nabla \psi - \nabla \psi \phi)$ is solenoidal. [4]

- c) Use Simpson's 1/3rd rule to find [4]

$\int_0^{0.6} e^{-x^2} dx$ by taking seven ordinates

- Q5) a) Find the work done by $\vec{F} = (2x + y^2) \hat{i} + (3y - 4x) \hat{j}$ in taking a particle around the parabolic arc $y = x^2$ from (0,0) to (1,1). [4]

- b) Apply Stokes's theorem to evaluate $\oint_C (4y dx + 2z dy + 6y dz)$ where C is curve of intersection of $x^2 + y^2 + z^2 = 2z$ and $z = x + 1$. [5]

- c) Evaluate $\iint_S (2y \hat{i} + yz \hat{j} + 2xz \hat{k}) \cdot d\vec{s}$ over the surface of region bounded by $y=0, y=3, x=0, z=0, x+2z=6$. [4]

OR

- Q6) a) Using Green's Lemma, evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F} = 3y \hat{i} + 2x \hat{j}$ and C is boundary of region bounded by $y=0, y=\sin x$ for $0 \leq x \leq \pi$ [4]

- b) Evaluate [5]

$$\iiint_s (z^2 - x) dy dz - xy dx dz + 3z dx dy$$

where s is closed surface of region bounded by $x=0, x=3, z=0, z=4-y^2$

c) Show that $\bar{E} = -\nabla\phi - 1/c \frac{\partial \bar{A}}{\partial t}$; $\bar{H} = \nabla \times \bar{A}$ are solutions of Maxwell's equations [4]

i) $\nabla \cdot \bar{H} = 0$ ii) $\nabla \times \bar{H} = 1/c \frac{\partial \bar{E}}{\partial t}$ if $\nabla \cdot \bar{A} + 1/c \frac{\partial \phi}{\partial t}$
and $\nabla^2 \bar{A} = 1/c^2 \frac{\partial^2 A}{\partial t^2}$

Q7) a) Show that the analytic function with constant amplitude is constant. [4]

b) By using Cauchy's integral formula, evaluate $\oint_c 2z^2 + z / z^2 - 1 dz$ [5]

Where c is the circle $|z-1| = 1$

c) Find the bilinear transformation which maps the points $z = -1, 0, 1$ on the points [4]

$w = 0, i, 3i$ of w-plane.

OR

Q8) a) If $u = \cos hx \cos y$ then find the harmonic conjugate v such that $f(z) = u + iv$ is analytical function. [4]

b) Evaluate [5]

$\int_c \frac{12z-7}{(z-1)^2(2z+3)} dz$ where c is the circle $|z|=2$ using Cauchy's residue theorem.

c) Show that the transformation $w = z + 1/z - 2i$ maps the circle $|z|=2$ into an ellipse. [4]

stopidsid.com